



Riparian forestry management and adult stream insects

R. A. Briers, J. H. R. Gee

► To cite this version:

R. A. Briers, J. H. R. Gee. Riparian forestry management and adult stream insects. Hydrology and Earth System Sciences Discussions, 2004, 8 (3), pp.545-549. hal-00304942

HAL Id: hal-00304942

<https://hal.science/hal-00304942>

Submitted on 1 Jan 2004

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Riparian forestry management and adult stream insects

Robert A. Briers and John H.R. Gee

Institute of Biological Sciences, University of Wales Aberystwyth, Edward Llwyd Building, Aberystwyth, Ceredigion SY23 3DA, UK

Email for corresponding author: r.briers@napier.ac.uk

Abstract

The impacts of coniferous plantation forestry on the biology of upland streams in the UK are firmly established. Whilst benthic communities have been well studied, very little research has considered the impacts of riparian forestry management on adult stream insects, yet the essentially terrestrial adult (reproductive) phase may be important in determining the abundance and distribution of larval stages. Riparian vegetation has a potentially strong impact on survival and success of adult stages through alteration of microclimate, habitat structure and potential food sources, in addition to effects carried over from larval stages. Here, current riparian management strategies are analysed in the light of available information on the ecology of adult stream insects. On the whole, management practices appear to favour adult stream insects, although an increase in tree cover in riparian areas could be beneficial, by providing more favourable microclimatic conditions for adults. This conclusion is drawn based on rather limited information, and the need for further research into the effects of riparian forestry management on adult stream insects is highlighted.

Keywords: microclimate, plantation, life history, riparian vegetation

Introduction

Since the beginning of the twentieth century, but particularly in the last 60 years, upland areas of Great Britain, particularly Wales and Scotland, have undergone extensive afforestation with exotic coniferous plantation (Stoner and Gee, 1985). Consequently, the catchments of many upland streams are heavily forested. Afforestation of riparian areas exacerbated the effects of acidification that was already occurring in the UK uplands (UKAWRG, 1988) and contributed to the deterioration of upland stream water quality (Harriman and Morrison, 1982; Neal *et al.*, 1992). Research into the effects of riparian forestry on invertebrate communities has focused almost exclusively on the benthic larval stages, notably the decline in primary productivity of streams due to channel shading (Winterbourn *et al.*, 1985) and alteration in physicochemical conditions and food resources, which result in changes in the diversity and abundance of benthic communities, due to the loss of taxa that are acid sensitive or herbivorous (Ormerod *et al.*, 1987, 1989, 1993; Gee and Smith, 1997). Management advice for riparian forestry areas has been focused similarly (Weatherley *et al.*, 1993; Forestry

Commission, 2000; Broadmeadow and Nisbet, 2002). No attention has been paid to the role of adult stages of aquatic insects in determining the distribution and abundance of populations. Many of the taxa that are numerically abundant in upland streams, notably Plecoptera, Trichoptera and Diptera, have adult stages which are effectively terrestrial, and which spend their lives in the riparian zone (Collier and Smith, 1998; Petersen *et al.*, 1999; Delettre and Morvan, 2000; Briers *et al.*, 2002). Mortality of adult stages in the terrestrial environment is high, in the region of 90% (Jackson and Fisher, 1986; Werneke and Zwick, 1992; Enders and Wagner, 1996), and hence factors influencing the distribution, survival and reproductive success of adults may be important in determining the abundance and distribution of larval stages (Statzner, 1977; Zwick, 1990; Timm, 1994; Harrison and Hildrew, 1998).

Adult stream insect ecology has only recently become an area of sustained research interest and, hence, information on the potential impacts of riparian forestry management is scattered and patchy. Firstly, relevant aspects of adult stream insect ecology have been reviewed, in conjunction with data

obtained during studies (Briers *et al.*, 2002; Briers *et al.*, 2003) of forested, clear-cut and moorland upland streams in the Plynlimon experimental catchment area (Hudson *et al.*, 1997). Current principles for the management of riparian forestry areas in the UK (Forestry Commission, 2000; Broadmeadow and Nisbet, 2002) are then analysed to determine whether management could be improved and areas for future study are highlighted.

From larva to adult — growing up with issues

Whilst research on the impacts of riparian forestry on stream insects has focused mainly on larvae rather than adults, much of the work also has relevance to adult stream insect ecology. Changes in larval food sources and stream physicochemical conditions associated with afforestation of riparian areas influence the size of adults at emergence indirectly through reductions in larval growth rates (Sweeney, 1984; Weatherley and Ormerod, 1990; Sweeney, 1993; Thomsen and Friberg, 2002). The smaller size at metamorphosis achieved by species in forested streams is linked to reductions in adult longevity, fecundity and reproductive success (Peterssen, 1987; Flecker *et al.*, 1988; Taylor *et al.*, 1998). Adult longevity is particularly important for species of stoneflies and caddisflies that feed as adults and spend longer periods in a terrestrial phase (Hynes, 1942; Petersson and Hasselrot, 1994; De Figueroa and Sanchez-Ortega, 2000).

In addition to influencing adult size at metamorphosis, alteration in developmental rates due to changes in food and stream temperature affects the timing of emergence. A comparison of the emergence dates of species of Plecoptera common to the clear felled Afon Hore and afforested Afon Hafren in mid-Wales, which differ in temperature (Crisp, 1997) and organic matter availability (Davies, 1996), [see Briers *et al.*, (2002) for details of the sites and species involved] showed a significant difference between the sites (paired t-test, $t_8 = 2.60$, $P = 0.032$), with species emerging on average 13 days later in the forested Hafren. Changes in the timing of emergence periods may have effects on the suitability of conditions for mating or flight and the availability of food sources, aspects of adult ecology that are covered in more detail subsequently.

Riparian microclimate

Microclimatic conditions experienced during the adult phase have an impact on survival and longevity. Air temperature and relative humidity are of particular importance, with higher temperatures and lower humidity reducing the adult

lifespan of aquatic groups (Jackson, 1988; Collier and Smith, 2000). Riparian vegetation has a key role in determining riparian microclimatic conditions. Conditions within mature forests with dense canopy cover are generally cooler with lower wind speeds and higher relative humidity, compared to areas outside the forest, such as clear cut or pasture (Chen *et al.*, 1993; Collier and Smith, 2000). The generally hotter, less humid and more variable conditions outside forestry could reduce adult insect lifespans. In a laboratory study of three New Zealand stoneflies, Collier and Smith (2000) found that the 96 hour LT_{50} (temperature at which 50% of individuals died over 96 hours) was exceeded for 25% of the time in a pasture riparian zone, compared to 0.1% of the time in a native forest catchment over two weeks during the emergence period.

Microclimate also influences the flight activity of aquatic species (Brindle, 1957; Waringer, 1991; Peng *et al.*, 1992a; Briers *et al.*, 2003). In common with terrestrial groups (e.g. Williams, 1961) flight activity of aquatic adults appears to be affected primarily by air temperature, but wind speed and humidity also influence flight in some groups. Many aquatic species, particularly Ephemeroptera (Savolainen, 1978), Diptera (Downes, 1969) and Trichoptera (Peterssen, 1989) form mating swarms. Swarming behaviour is inhibited at low temperatures (Solem, 1976; Savolainen, 1978; Peterssen, 1989; Jackson and Resh, 1991), therefore the lower temperatures characteristic of dense riparian forestry may restrict opportunities for species to swarm and mate.

Habitat structure

The extent and nature of riparian forestry alters the structure of the riparian habitat. Some studies have suggested that dense riparian vegetation may restrict the dispersal of individuals away from the stream channel and, hence, influence the likelihood of colonisation or transfer of individuals between neighbouring sites (Jackson and Resh, 1989; Petersen *et al.*, 1999). Studies of stonefly dispersal at moorland, clear-cut and forested streams in mid Wales (Briers *et al.*, 2002) found that the rate of decline in numbers with distance from the streams did not vary with riparian land-use. However, in a lowland agricultural landscape in France there was a clear effect of habitat structure on the dispersal of Chironomidae (Delettre and Morvan, 2000). The differences between the two studies in responses to variation in riparian vegetation may reflect variation in adult behaviour. Chironomids (and other swarming species) utilize trees and other vegetation as markers for mating swarms (Downes, 1969; Savolainen *et al.*, 1993), whereas stoneflies do not swarm. Variation in riparian vegetation structure may also alter the predation risk faced by adult insects emerging

from streams. Riparian areas support an abundance of predators, including birds, lizards and spiders, that exploit adult aquatic insects (Jackson and Fisher, 1986; Machtans *et al.*, 1996; Sabo and Power, 2002; Sanzone *et al.*, 2003) and may contribute significantly to the mortality of adult stages.

Food sources

Ephemeroptera, Chironomidae and most Trichoptera are not known to feed in their adult phase, but many species of stoneflies (Hynes, 1942, 1976; De Figueroa and Sanchez-Ortega, 2000; K.H. Macneale, *pers. comm.*) and some caddisflies (Petersson and Hasselrot, 1994), require to feed to mate and mature eggs successfully. There is very limited information on the terrestrial nutritional requirements of species, but pollen, lichen, fungi and algae are common components of adult stonefly diets (Hynes, 1942; De Figueroa and Sanchez-Ortega, 2000; Smith and Collier, 2000) and hence multiple sources of food may be important. Differences in food supplies between riparian vegetation types may therefore influence longevity and egg production of stonefly species (Sweeney, 1993; Smith and Collier, 2000).

Current riparian forestry management in relation to adult stream insects

Guidelines for the management of riparian areas of plantation forestry in the UK (Forestry Commission, 1990, 1991, 2000; see also Broadmeadow and Nisbet, 2002) have focused on the establishment and maintenance of 'buffer strips' – uncultivated areas whose primary function is to protect the stream channel and riparian zone from acidic drainage and sediment from adjacent forestry (Forestry Commission, 2000). The recommended size of the buffer strip on each bank varies between 5 and 20 m depending on the size of the stream channel. Management of buffer strips is aimed at maintaining open/partially wooded (primarily broadleaf) conditions. Trees providing heavy shade are to be used only sparsely and not within 10 m of the stream channel to prevent excessive shading and encourage the growth of ground vegetation to trap sediment from adjacent forestry (Forestry Commission, 2000).

The majority of adult aquatic insects do not disperse a significant distance away from the stream channel. Typical dispersal distance may vary with taxon, but the decline in numbers with increasing distance from the channel has been commonly represented as a negative exponential or inverse power function (Griffith *et al.*, 1998; Petersen *et al.*, 1999; Briers *et al.*, 2002). For this reason, it is the areas

immediately adjacent to the stream channel that are likely to be most important to the majority of the adult stream insect population.

The maintenance of a diverse range of riparian vegetation as currently recommended is likely to be beneficial to adult stream insects. Tree cover will provide more suitable microclimatic conditions, food sources and shelter from predation. Isolated individual or small groups of trees can play an important role as markers for swarming species (Peng *et al.*, 1992b). However, the extent to which variation in microclimatic conditions is reduced within forestry relative to open ground is highly dependent on the size of the forest patch. Relatively small patches of trees suffer from strong microclimatic edge effects (Chen *et al.*, 1995; Brososke *et al.*, 1997) and, hence, more extensive riparian tree cover may be necessary to provide suitable microclimatic conditions.

The location of riparian forestry should also be considered. While it is clearly undesirable to maintain highly shaded conditions over the length of the stream channel, with the limited dispersal of some taxa [90% of adult stoneflies travelled less than 11 m away from upland streams in Wales, regardless of riparian vegetation type (Briers *et al.*, 2002)] it is important to maintain some forestry immediately adjacent to the channel. If such patches are located some distance away from the stream channel, the greater dispersal distances required to reach them will increase the risk of predation or desiccation.

Areas for future study

Information on the ecology of adult stream insects is far from complete, so the conclusions that can be drawn on the potential impacts of riparian forestry management are limited. Of a number of areas where further information is required, two are explored here.

ADULT VERSUS LARVAL REQUIREMENTS

A recent study (Winterbourn and Crowe, 2001) has shown that some species of adult aquatic insect show significant directionality of flight relative to the location of riparian forest patches. Winterbourn and Crowe (2001) interpreted this as a function of larval habitat requirements, but areas suitable for larvae may not necessarily coincide with areas favourable for adult survival and this may lead to differences in the distribution patterns of different life stages of the same species. Some work is already underway exploring this area (S. Flint and M. Dobson, unpublished data), but this is clearly an area where further research would be beneficial to determine the relative contribution of adult and larval life

stages to small scale abundance and distribution patterns relative to riparian vegetation.

ADULT FOOD SOURCES

The significance of terrestrial feeding for adult stoneflies and some caddisflies, which are often abundant in forested upland streams, is well established. However, the implications of variation in adult food supply in different riparian vegetation types for the reproductive success of species is far from clear. Location of appropriate food sources is likely to be an important factor in adult habitat selection (Harper, 1973) and further information on adult requirements would contribute to management of riparian forestry.

Conclusions

Despite the potential importance of adult success in determining the abundance and distribution of benthic larval stages, the adult phase of aquatic insect lifecycles has been neglected when considering the impact of riparian forestry management. Riparian areas in forestry have a number of roles and management of these areas will ultimately represent a trade-off between sometimes conflicting requirements of the different functions. An evaluation of riparian management policies in the light of available information on the factors influencing adult aquatic insect success suggests that current practices are generally beneficial to adult aquatic insects, although a greater emphasis on areas of riparian forest cover, more in keeping with policies in the United States e.g. Palone and Todd (1997) might well improve the success of adult aquatic insects by providing a more favourable microclimate, improving adult survival. However, there is clearly a need for further research to evaluate the effects of riparian forestry management on adult stream insects.

Acknowledgements

The authors are grateful to James Laing, Conservation & Environment Support Officer, Forest Enterprise Wales for providing information on current management practices. This work was supported by grant GR3/12114 from the UK Natural Environment Research Council.

References

Briers, R.A., Cariss, H.M. and Gee, J.H.R., 2002. Dispersal of adult stoneflies (Plecoptera) from upland streams draining catchments with contrasting land-use. *Arch. Hydrobiol.*, **155**, 627–644.

- Briers, R.A., Cariss, H.M. and Gee, J.H.R., 2003. Flight activity of adult stoneflies in relation to weather. *Ecol. Entomol.*, **28**, 31–40.
- Brindle, A., 1957. The effect of temperature and humidity on the flight of Trichoptera. *Entomol. Mon. Mag.*, **93**, 63–66.
- Broadmeadow, S. and Nisbet, T., eds. (2002) *The effect of riparian forest management on the freshwater environment*. SNIFFER, Edinburgh, UK.
- Brososke, K.D., Chen, J.Q., Naiman, R.J. and Franklin, J.F., 1997. Harvesting effects on microclimatic gradients from small streams to uplands in western Washington. *Ecol. Appl.*, **7**, 1188–1200.
- Chen, J.Q., Franklin, J.F. and Spies, T.A., 1993. Contrasting microclimates among clear-cut, edge, and interior of old-growth Douglas fir forest. *Agr. Forest Meteorol.*, **63**, 219–237.
- Chen, J.Q., Franklin, J.F. and Spies, T.A., 1995. Growing season microclimatic gradients from forest clear-cut edges into old-growth Douglas fir forests. *Ecol. Appl.*, **5**, 74–86.
- Collier, K.J. and Smith, B.J., 1998. Dispersal of adult caddisflies (Trichoptera) into forests alongside three New Zealand streams. *Hydrobiol.*, **361**, 53–65.
- Collier, K.J. and Smith, B.J., 2000. Interactions of adult stoneflies (Plecoptera) with riparian zones I. effects of air temperature and humidity on longevity. *Aquat. Insects*, **22**, 275–284.
- Crisp, D.T., 1997. Water temperature of Plynlimon streams. *Hydrol. Earth Syst. Sci.*, **1**, 535–540.
- Davies, A.L., 1996. *Organic matter availability in upland streams: spatial and temporal patterns in relation to land-use*. Ph.D Thesis, University of Wales, Aberystwyth, UK.
- De Figueroa, J.M.T. and Sanchez-Ortega, A., 2000. Imaginal feeding of twelve Nemouroidean stonefly species (Insecta, Plecoptera). *Ann. Entomol. Soc. Amer.*, **93**, 251–253.
- Delettre, Y.R. and Morvan, N., 2000. Dispersal of adult aquatic chironomidae (Diptera) in agricultural landscapes. *Freshwater Biol.*, **44**, 399–411.
- Downes, J.A., 1969. The swarming and mating flight of Diptera. *Ann. Rev. Entomol.*, **14**, 271–298.
- Enders, G. and Wagner, R., 1996. Mortality of *Apatania fimbriata* (Insecta: Trichoptera) during embryonic, larval and adult life stages. *Freshwater Biol.*, **36**, 93–104.
- Flecker, A.S., Allan, J.D. and McClintock, N.L., 1988. Male body size and mating success in swarms of the mayfly *Epeorus longimanus*. *Holarctic Ecol.*, **11**, 280–285.
- Forestry Commission, 1990. *Forest Nature Conservation Guidelines*. HMSO, London, UK.
- Forestry Commission, 1991. *Forests and Water Guidelines*, 2nd edn. HMSO, London, UK.
- Forestry Commission, 2000. *Forests and Water Guidelines*, 4th edn. The Forestry Commission, Edinburgh, UK.
- Gee, J.H.R. and Smith, B.D., 1997. Benthic invertebrates in the headwaters of the Wye and Severn: effects of forestry and clear-felling. *Hydrol. Earth Syst. Sci.*, **1**, 549–556.
- Griffith, M.B., Barrows, E.M. and Perry, S.A., 1998. Lateral dispersal of adult aquatic insects (Plecoptera, Trichoptera) following emergence from headwater streams in forested Appalachian catchments. *Ann. Entomol. Soc. Amer.*, **91**, 195–201.
- Harper, P.P., 1973. Life histories of Nemoridae and Leuctridae in Southern Ontario (Plecoptera). *Hydrobiol.*, **41**, 309–356.
- Harriman, R. and Morrison, B.R.S., 1982. Ecology of streams draining forested and non-forested catchments in an area of central Scotland subject to acid precipitation. *Hydrobiol.*, **88**, 251–263.
- Harrison, S.S.C. and Hildrew, A.G., 1998. Distribution dynamics of epilithic insects in a lake littoral. *Arch. Hydrobiol.*, **143**, 275–293.

- Hudson, J.A., Gilman, K. and Calder, I.R., 1997. Land use and water issues in the uplands with reference to the Plynlimon study. *Hydrol. Earth Syst. Sci.*, **1**, 389–397.
- Hynes, H.B.N., 1942. A study of feeding in adult stoneflies (Plecoptera). *Proc. R. Entomol. Soc. Lond. (A)*, **17**, 81–82.
- Hynes, H.B.N., 1976. Biology of Plecoptera. *Ann. Rev. Entomol.*, **21**, 135–153.
- Jackson, J.K., 1988. Diel emergence, swarming and longevity of selected adult aquatic insects from a Sonoran Desert stream. *Amer. Midland Naturalist*, **119**, 344–352.
- Jackson, J.K. and Fisher, S.G., 1986. Secondary production, emergence, and export of aquatic insects of a Sonoran desert stream. *Ecol.*, **67**, 629–638.
- Jackson, J.K. and Resh, V.H., 1989. Distribution and abundance of adult aquatic insects in the forest adjacent to a northern California stream. *Environ. Entomol.*, **18**, 278–283.
- Jackson, J.K. and Resh, V.H., 1991. Periodicity in mate attraction and flight activity of three species of caddisflies (Trichoptera). *J. N. Amer. Benth. Soc.*, **10**, 198–209.
- Machtans, C.S., Villard, M.-A. and Hannon, S.J., 1996. Use of riparian buffer strips as movement corridors by forest birds. *Conserv. Biol.*, **10**, 1366–1379.
- Neal, C., Smith, C.J. and Hill, S., (Eds.), 1992. *Forestry impact on upland water quality*. Institute of Hydrology, Wallingford, UK.
- Ormerod, S.J., Donald, A.P. and Brown, S.J., 1989. The influence of plantation forestry on the pH and aluminium concentration of upland Welsh streams: a re-examination. *Environ. Poll.*, **62**, 47–63.
- Ormerod, S.J., Mawle, G.W. and Edwards, R.W., 1987. The influence of forest on aquatic fauna. In: *Environmental aspects of plantation forestry in Wales*, J.E.G. Good (Ed.), Institute of Terrestrial Ecology, Grange-Over-Sands, UK. 37–49.
- Ormerod, S.J., Rundle, S.D., Lloyd, E.C. and Douglas, A.A., 1993. The influence of riparian management on the habitat structure and macroinvertebrate communities of upland streams draining plantation forest. *J. Appl. Ecol.*, **30**, 13–24.
- Palone, R.S. and Todd, A.H., (Eds.), 1997. *Chesapeake Bay riparian handbook: a guide for establishing and maintaining riparian forest buffers*. USDA Forest Service, Radnor, PA, USA.
- Peng, R.K., Fletcher, C.R. and Sutton, S.L., 1992a. The effect of microclimate on flying dipterans. *Int. J. Biometeorol.*, **36**, 69–76.
- Peng, R.K., Sutton, S.L. and Fletcher, C.R., 1992b. Spatial and temporal distribution patterns of flying Diptera. *J. Zool. (Lond.)*, **228**, 329–340.
- Petersen, I., Winterbottom, J.H., Orton, S., Friberg, N., Hildrew, A.G., Spiers, D.C. and Gurney, W.S.C., 1999. Emergence and lateral dispersal of adult Plecoptera and Trichoptera from Broadstone Stream, U.K. *Freshwater Biol.*, **42**, 401–416.
- Petersen, E., 1987. Weight-associated male mating success in the swarming caddis fly, *Mystacides azureus* L. *Ann. Zool. Fenn.*, **24**, 335–339.
- Petersen, E., 1989. Swarming activity patterns and seasonal decline in adult size in some caddisflies (Trichoptera: Leptoceridae). *Aquat. Insect.*, **11**, 17–28.
- Petersen, E. and Hasselrot, A.T., 1994. Mating and nectar feeding in the psychomid caddisfly *Tinodes waeneri*. *Aquat. Insect.*, **16**, 177–187.
- Sabo, J.L. and Power, M.E., 2002. River-watershed exchange: effects of riverine subsidies on riparian lizards and their terrestrial prey. *Ecol.*, **83**, 1860–1869.
- Sanzone, D.M., Meyer, J.L., Marti, E., Gardiner, E.P., Tank, J.L. and Grimm, N.B., 2003. Carbon and nitrogen transfer from a desert stream to riparian predators. *Oecologia*, **134**, 238–250.
- Savolainen, E., 1978. Swarming in Ephemeroptera: the mechanism of swarming and the effects of illumination and weather. *Ann. Zool. Fenn.*, **15**, 17–52.
- Savolainen, E., Saura, A. and Hantula, J., 1993. Mode of swarming in relation to reproductive isolation in mayflies. *Evolution*, **47**, 1796–1804.
- Smith, B.J. and Collier, K.J., 2000. Interactions of adult stoneflies (Plecoptera) with riparian zones II. diet. *Aquat. Insect.*, **22**, 285–296.
- Solem, J.O., 1976. Studies on the behaviour of adults of *Phryganea bipunctata* and *Agrypnia obsoleta* (Trichoptera). *Norweg. J. Entomol.*, **23**, 23–28.
- Statzner, B., 1977. The effects of flight behaviour on the larval abundance of Trichoptera in the Schierenseebrooks (North Germany). In: *Proc. Second Int. Symp. on Trichoptera*, M.I. Crichton (Ed.), Junk, The Hague, Reading. 121–134.
- Stoner, J.H. and Gee, A.S., 1985. Effects of forestry on water quality and fish in Welsh rivers and lakes. *J. Inst. Water Eng. Sci.*, **39**, 27–45.
- Sweeney, B.W., 1984. Factors influencing life-history patterns of aquatic insects. In: *The Ecology of Aquatic Insects*. V.H. Resh and D.M. Rosenberg (Eds.), Praeger Scientific, New York, USA. 56–100.
- Sweeney, B.W., 1993. Effects of streamside vegetation on macroinvertebrate communities of White Clay Creek in Eastern North America. *Proc. Acad. Natur. Sci. Phil. A*, **144**, 291–340.
- Taylor, B.W., Anderson, C.R. and Peckarsky, B.L., 1998. Effects of size at metamorphosis on stonefly fecundity, longevity, and reproductive success. *Oecologia*, **114**, 494–502.
- Thomsen, A.G. and Friberg, N., 2002. Growth and emergence of the stonefly *Leuctra nigra* in coniferous streams with contrasting pH. *Freshwater Biol.*, **47**, 1159–1172.
- Timm, T., 1994. Reasons for the shift in dominance between *Simulium* (N.) *vernum* and *Simulium* (S.) *ornatum* (Diptera: Simuliidae) along the continuum of an unpolluted lowland stream. *Arch. Hydrobiol.*, **131**, 199–210.
- UKAWRG, 1988. *Acidity in United Kingdom Fresh Waters*, HMSO, London, UK.
- Waringer, J.A., 1991. Phenology and the influence of meteorological parameters on the catching success of light-trapping for Trichoptera. *Freshwater Biol.*, **25**, 307–319.
- Weatherley, N.S., Lloyd, E.C., Rundle, S.D. and Ormerod, S.J., 1993. Management of conifer plantations for the conservation of stream macroinvertebrates. *Biol. Conserv.*, **63**, 171–176.
- Weatherley, N.S. and Ormerod, S.J., 1990. Forests and the temperature of upland streams in Wales: a modelling exploration of the biological effects. *Freshwater Biol.*, **24**, 109–122.
- Werneke, U. and Zwick, P., 1992. Mortality of the terrestrial adult and aquatic nymphal life stages of *Baetis vernus* and *Baetis rhodani* in the Breitenbach, Germany (Insecta: Ephemeroptera). *Freshwater Biol.*, **28**, 249–255.
- Williams, C.B., 1961. Studies in the effect of weather conditions on the activity and abundance of insect populations. *Phil. Trans. Roy. Soc. Lond. (B)*, **244**, 331–378.
- Winterbourn, M.J. and Crowe, A.L.M., 2001. Flight activity of insects along a mountain stream: is directional flight adaptive? *Freshwater Biol.*, **46**, 1479–1489.
- Winterbourn, M.J., Hildrew, A.G. and Box, A., 1985. Structure and grazing of stone surface organic layers in some acid streams of southern England. *Freshwater Biol.*, **15**, 363–374.
- Zwick, P., 1990. Emergence, maturation and upstream oviposition flights of Plecoptera from the Breitenbach, with notes on the adult phase as a possible control of stream insect populations. *Hydrobiol.*, **194**, 207–233.